

## DEEPLY VIRTUAL COMPTON SCATTERING AND GENERALIZED PARTON DISTRIBUTIONS

*A. Flachi*<sup>1</sup>, *K. Gudima*<sup>2</sup>, *L.L. Jenkovszky*<sup>3</sup>,  
*V. Kuvshinov*<sup>4</sup>, *V.K. Magas*<sup>3,5</sup>

<sup>1</sup>Universidad Autonoma de Barselona  
(*08193 Bellaterra, Barselona, Spain*),

<sup>2</sup>Institute of Applied Physics, Acad. of Sci. of Moldova  
(*5, Academy Str., Kishinev, MD-2028, Moldova*),

<sup>3</sup>Bogolyubov Institute for Theoretical Physics,  
Nat. Acad. Sci. of Ukraine  
(*14b, Metrolohichna Str., 03143 Kyiv, Ukraine*),

<sup>4</sup>Institute of Physics, Belarus Nat. Acad. of Sci.  
(*68, F.Skarina Ave., 220072 Minsk,*  
*Republik of Belarus*),

<sup>5</sup>Physics Department, Instituto Superior Tecnico  
(*Rovisco Pais Ave., 1049-001 Lisbon, Portugal*)

### S u m m a r y

An off-mass-shell continuation of the dual model with Mandelstam analyticity is proposed. It can serve as a prototype model for deeply virtual Compton scattering (DVCS) and generalized parton distributions (GPD). With the spin dependence included, the model is fitted to the data on inelastic electron-proton scattering.